

- [Return to BCAlgTori](#)

FlabbyResolutionBC.gap

Definition of M_G

Let G be a finite subgroup of $\mathrm{GL}(n, \mathbb{Z})$. The G -lattice M_G of rank n is defined to be the G -lattice with a \mathbb{Z} -basis $\{u_1, \dots, u_n\}$ on which G acts by

$$\sigma(u_i) = \sum_{j=1}^n a_{i,j} u_j \quad (1)$$

for any $\sigma = [a_{i,j}] \in G$.

Hminus1

▶ Hminus1(G)

returns the Tate cohomology group $\widehat{H}^{-1}(G, M_G)$ for a finite subgroup $G \leq \mathrm{GL}(n, \mathbb{Z})$.

H0

▶ H0(G)

returns the Tate cohomology group $\widehat{H}^0(G, M_G)$ for a finite subgroup $G \leq \mathrm{GL}(n, \mathbb{Z})$.

H1

▶ H1(G)

returns the cohomology group $H^1(G, M_G)$ for a finite subgroup $G \leq \mathrm{GL}(n, \mathbb{Z})$.

Sha1Omega

▶ Sha1Omega(G)

returns $Sha_w^1(G, M_G)$.

Sha1OmegaTr

▶ Sha1OmegaTr(G)

returns $Sha_w^1(G, (M_G)^\circ)$.

ShaOmega

▸ ShaOmega(G, n)

returns $Sha_w^n(G, M_G)$ for G -lattice M_G .
This function needs HAP package in GAP.

ShaOmegaFromGroup

▸ ShaOmegaFromGroup(M, n, G)

returns $Sha_w^n(G, M)$ for G -lattice M .
This function needs HAP package in GAP.

TorusInvariants

▸ TorusInvariants(G)

returns $TI_G = [l_1, l_2, l_3, l_4]$ where

$$l_1 = \begin{cases} 0 & \text{if } [M_G]^{fl} = 0, \\ 1 & \text{if } [M_G]^{fl} \neq 0 \text{ but is invertible,} \\ 2 & \text{if } [M_G]^{fl} \text{ is not invertible,} \end{cases}$$

$$\begin{aligned} l_2 &= H^1(G, [M_G]^{fl}) \simeq Sha_w^1(G, [M_G]^{fl}), \\ l_3 &= Sha_w^1(G, (M_G)^\circ) \simeq Sha_w^2(G, ([M_G]^{fl})^\circ), \\ l_4 &= H^1(G, ([M_G]^{fl})^{fl}) \simeq Sha_w^2(G, [M_G]^{fl}) \text{ via the command H1}(\mathbf{G}). \end{aligned}$$

TorusInvariantsHAP

▸ TorusInvariantsHap(G)

returns $TI_G = [l_1, l_2, l_3, l_4]$ where

$$l_1 = \begin{cases} 0 & \text{if } [M_G]^{fl} = 0, \\ 1 & \text{if } [M_G]^{fl} \neq 0 \text{ but is invertible,} \\ 2 & \text{if } [M_G]^{fl} \text{ is not invertible,} \end{cases}$$

$$\begin{aligned} l_2 &= H^1(G, [M_G]^{fl}) \simeq Sha_w^1(G, [M_G]^{fl}), \\ l_3 &= Sha_w^1(G, (M_G)^\circ) \simeq Sha_w^2(G, ([M_G]^{fl})^\circ), \\ l_4 &= Sha_w^2(G, [M_G]^{fl}) \text{ via the command ShaOmegaFromGroup}([M_G]^{fl}, 2, G). \end{aligned}$$

This function needs HAP package in GAP.

ConjugacyClassesSubgroups2TorusInvariants

▸ `ConjugacyClassesSubgroups2TorusInvariants(G)`

returns the records `ConjugacyClassesSubgroups2` and `TorusInvariants` where `ConjugacyClassesSubgroups2` is the list $[g_1, \dots, g_m]$ of conjugacy classes of subgroups of $G \leq \text{GL}(n, \mathbb{Z})$ with the fixed ordering via the function `ConjugacyClassesSubgroups2(G)` ([HY17, Section 4.1]) and `TorusInvariants` is the list $[\text{TorusInvariants}(g_1), \dots, \text{TorusInvariants}(g_m)]$ via the function `TorusInvariants(G)`.

PossibilityOfStablyEquivalentSubdirectProducts

▸ `PossibilityOfStablyEquivalentSubdirectProducts(G, G' ,
ConjugacyClassesSubgroups2TorusInvariants(G),
ConjugacyClassesSubgroups2TorusInvariants(G'))`

returns the list l of the subdirect products $\widetilde{H} \leq G \times G'$ of G and G' up to $(\text{GL}(n_1, \mathbb{Z}) \times \text{GL}(n_2, \mathbb{Z}))$ -conjugacy which satisfy $TI_{\varphi_1(H)} = TI_{\varphi_2(H)}$ for any $H \leq \widetilde{H}$ where $\widetilde{H} \leq G \times G'$ is a subdirect product of G and G' which acts on M_G and $M_{G'}$ through the surjections $\varphi_1 : \widetilde{H} \rightarrow G$ and $\varphi_2 : \widetilde{H} \rightarrow G'$ respectively (indeed, this function computes it for H up to conjugacy for the sake of saving time). In particular, if the length of the list l is zero, then we find that $[M_G]^{fl}$ and $[M_{G'}]^{fl}$ are not weak stably k -equivalent.

FlabbyResolutionLowRank

▸ `FlabbyResolutionLowRank(G).actionF`

returns the matrix representation of the action of G on F where F is a suitable flabby class of M_G ($F = [M_G]^{fl}$) with low rank by using backtracking techniques (see [HY17, Chapter 5], see also [HHY Algorithm 4.1 (3)]).

Each isomorphism class of irreducible permutation \widetilde{H} -lattices corresponds to a conjugacy class of subgroup H of \widetilde{H} by $H \leftrightarrow \mathbb{Z}[\widetilde{H}/H]$. Let $H_1 = \{1\}, \dots, H_r = \widetilde{H}$ be all conjugacy classes of subgroups of \widetilde{H} whose ordering corresponds to the GAP function `ConjugacyClassesSubgroups2(\widetilde{H})` (see [HY17, Section 4.1, page 42]).

We suppose that $[F] = [F']$ as \widetilde{H} -lattices. Then we have

$$\left(\bigoplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus x_i} \right) \oplus F^{\oplus b_1} \simeq \left(\bigoplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus y_i} \right) \oplus F'^{\oplus b_1} \quad (2)$$

where $b_1 = 1$. We write the equation (2) as

$$\bigoplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a_i} \simeq (F - F')^{\oplus (-b_1)} \quad (3)$$

formally where $a_i = x_i - y_i \in \mathbb{Z}$. Then we may consider " $F - F'$ " formally in the sene of (2). By computing some $\text{GL}(n, \mathbb{Z})$ -conjugacy class invariants, we will give a necessary condition for $[F] = [F']$.

Let $\{c_1, \dots, c_r\}$ be a set of complete representatives of the conjugacy classes of \widetilde{H} . Let $A_i(c_j)$ be the matrix representation of the factor coset action of $c_j \in \widetilde{H}$ on $\mathbb{Z}[\widetilde{H}/H_i]$ and $B(c_j)$ be the matrix representation of the action of $c_j \in \widetilde{H}$ on $F - F'$.

By (3), for each $c_j \in \widetilde{H}$, we have

$$\sum_{i=1}^r a_i \text{tr } A_i(c_j) + b_1 \text{tr } B(c_j) = 0 \quad (4)$$

where $\text{tr } A$ is the trace of the matrix A . Similarly, we consider the rank of $H^0 = \widehat{Z}^0$. For each H_j , we get

$$\sum_{i=1}^r a_i \text{rank } \widehat{Z}^0(H_j, \mathbb{Z}[\widetilde{H}/H_i]) + b_1 \text{rank } \widehat{Z}^0(H_j, F - F') = 0. \quad (5)$$

Finally, we compute \widehat{H}^0 . Let $\text{Sy}_p(A)$ be a p -Sylow subgroup of an abelian group A . $\text{Sy}_p(A)$ can be written as a direct product of cyclic groups uniquely. Let $n_{p,e}(\text{Sy}_p(A))$ be the number of direct summands of cyclic groups of order p^e . For each H_j, p, e , we get

$$\sum_{i=1}^r a_i n_{p,e}(\text{Sy}_p(\widehat{H}^0(H_j, \mathbb{Z}[\widetilde{H}/H_i]))) + b_1 n_{p,e}(\text{Sy}_p(\widehat{H}^0(H_j, F - F')))) = ($$

By the equalities (4), (5) and (6), we may get a system of linear equations in a_1, \dots, a_r, b_1 over \mathbb{Z} . Namely, we have that $[F] = [F']$ as \widetilde{H} -lattices \implies there exist $a_1, \dots, a_r \in \mathbb{Z}$ and $b_1 = \pm 1$ which satisfy (3) \implies this system of linear equations has an integer solution in a_1, \dots, a_r with $b_1 = \pm 1$.

In particular, if this system of linear equations has no integer solutions, then we conclude that $[F] \neq [F']$ as \widetilde{H} -lattices.

PossibilityOfStablyEquivalentFSubdirectProduct

▸ PossibilityOfStablyEquivalentFSubdirectProduct($H\sim$)

returns a basis $\mathcal{L} = \{l_1, \dots, l_s\}$ of the solution space $\{[a_1, \dots, a_r, b_1] \mid a_i, b_1 \in \mathbb{Z}\}$ of the system of linear equations which is obtained by the equalities (4), (5) and (6) and gives all possibilities that establish the equation (3) for a subdirect product $\widetilde{H} \leq G \times G'$ of G and G' .

PossibilityOfStablyEquivalentMSubdirectProduct

▸ PossibilityOfStablyEquivalentMSubdirectProduct($H\sim$)

returns the same as PossibilityOfStablyEquivalentFSubdirectProduct($H\sim$) but with respect to M_G and $M_{G'}$ instead of F and F' .

PossibilityOfStablyEquivalentFSubdirectProduct with "H2" option

▸ PossibilityOfStablyEquivalentFSubdirectProduct($H\sim$:H2)

returns the same as PossibilityOfStablyEquivalentFSubdirectProduct($H\sim$) but using also the additional equality

$$\sum_{i=1}^r a_i n_{p,e}(Sy_p(H^2(\widetilde{H}, \mathbb{Z}[\widetilde{H}/H_i]))) + b_1 n_{p,e}(Sy_p(H^2(\widetilde{H}, F - F')))) = 0$$

and the equalities (4), (5) and (6).

PossibilityOfStablyEquivalentMSubdirectProduct with "H2" option

▸ PossibilityOfStablyEquivalentMSubdirectProduct($H\sim$:H2)

returns the same as PossibilityOfStablyEquivalentFSubdirectProduct($H\sim$:H2) but with respect to M_G and $M_{G'}$ instead of F and F' .

In general, we will provide a method in order to confirm the isomorphism

$$\left(\bigoplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a_i} \right) \oplus F^{\oplus b_1} \simeq \left(\bigoplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a'_i} \right) \oplus F'^{\oplus b'_1} \quad (8)$$

with $a_i, a'_i \geq 0$, $b_1, b'_1 \geq 1$, although it is needed by trial and error.

Let G_1 (resp. G_2) be the matrix representation group of the action of \widetilde{H} on the left-hand side $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a_i}) \oplus F^{\oplus b_1}$ (resp. the right-hand side $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a'_i}) \oplus F'^{\oplus b'_1}$) of the isomorphism (8). Let $\mathcal{P} = \{P_1, \dots, P_m\}$ be a basis of the solution space of $G_1 P = P G_2$ where $m = \text{rank}_{\mathbb{Z}} \text{Hom}(G_1, G_2) = \text{rank}_{\mathbb{Z}} \text{Hom}_{\widetilde{H}}(M_{G_1}, M_{G_2})$. Our aim is to find the matrix P which satisfies $G_1 P = P G_2$ by using computer effectively. If we can get a matrix P with $\det P = \pm 1$, then G_1 and G_2 are $\text{GL}(n, \mathbb{Z})$ -conjugate where n is the rank of both sides of (8) and hence the isomorphism (8) established. This implies that the flabby class $[F^{\oplus b_1}] = [F'^{\oplus b'_1}]$ as \widetilde{H} -lattices.

StablyEquivalentFCheckPSubdirectProduct

▸ `StablyEquivalentFCheckPSubdirectProduct(\widetilde{H} , $\mathbf{l1}$, $\mathbf{l2}$)`

returns a basis $\mathcal{P} = \{P_1, \dots, P_m\}$ of the solution space of $G_1 P = P G_2$ where $m = \text{rank}_{\mathbb{Z}} \text{Hom}(G_1, G_2)$ and G_1 (resp. G_2) is the matrix representation group of the action of \widetilde{H} on $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a_i}) \oplus F^{\oplus b_1}$ (resp. $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a'_i}) \oplus F'^{\oplus b'_1}$) with the isomorphism (8) for a subdirect product $\widetilde{H} \leq G \times G'$ of G and G' , and lists $\mathbf{l1} = [a_1, \dots, a_r, b_1]$, $\mathbf{l2} = [a'_1, \dots, a'_r, b'_1]$, if P exists. If such P does not exist, this returns `[]`.

StablyEquivalentMCheckPSubdirectProduct

▸ `StablyEquivalentMCheckPSubdirectProduct(\widetilde{H} , $\mathbf{l1}$, $\mathbf{l2}$)`

returns the same as `StablyEquivalentFCheckPSubdirectProduct(\widetilde{H} , $\mathbf{l1}$, $\mathbf{l2}$)` but with respect to M_G and $M_{G'}$ instead of F and F' .

StablyEquivalentFCheckMatSubdirectProduct

▸ `StablyEquivalentFCheckMatSubdirectProduct(\widetilde{H} , $\mathbf{l1}$, $\mathbf{l2}$, P)`

returns true if $G_1 P = P G_2$ and $\det P = \pm 1$ where G_1 (resp. G_2) is the matrix representation group of the action of \widetilde{H} on $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a_i}) \oplus F^{\oplus b_1}$ (resp. $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a'_i}) \oplus F'^{\oplus b'_1}$) with the isomorphism (8) for a subdirect product $\widetilde{H} \leq G \times G'$ of G and G' , and lists $\mathbf{l1} = [a_1, \dots, a_r, b_1]$, $\mathbf{l2} = [a'_1, \dots, a'_r, b'_1]$. If not, this returns false.

StablyEquivalentMCheckMatSubdirectProduct

▸ `StablyEquivalentMCheckMatSubdirectProduct(\widetilde{H} , $\mathbf{l1}$, $\mathbf{l2}$, P)`

returns the same as `StablyEquivalentFCheckMatSubdirectProduct(\widetilde{H} , $\mathbf{l1}$, $\mathbf{l2}$, P)` but with respect to M_G and $M_{G'}$ instead of F and F' .

StablyEquivalentFCheckGenSubdirectProduct

▸ StablyEquivalentFCheckGenSubdirectProduct(\widetilde{H} , l_1, l_2)

returns the list $[\mathcal{M}_1, \mathcal{M}_2]$ where $\mathcal{M}_1 = [g_1, \dots, g_t]$ (resp. $\mathcal{M}_2 = [g'_1, \dots, g'_t]$) is a list of the generators of G_1 (resp. G_2) which is the matrix representation group of the action of \widetilde{H} on $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a_i}) \oplus F^{\oplus b_1}$ (resp. $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a'_i}) \oplus F'^{\oplus b'_1}$) with the isomorphism (8) for a subdirect product $\widetilde{H} \leq G \times G'$ of G and G' , and lists $l_1 = [a_1, \dots, a_r, b_1]$, $l_2 = [a'_1, \dots, a'_r, b'_1]$.

StablyEquivalentMCheckGenSubdirectProduct

▸ StablyEquivalentMCheckGenSubdirectProduct(\widetilde{H} , l_1, l_2)

returns the same as StablyEquivalentFCheckGenSubdirectProduct(\widetilde{H} , l_1, l_2) but with respect to M_G and $M_{G'}$ instead of F and F' .

By applying the function StablyEquivalentFCheckPSubdirectProduct, we get a basis $\mathcal{P} = \{P_1, \dots, P_m\}$ of the solution space of $G_1 P = P G_2$ with $\det P_i = \pm 1$ for some $1 \leq i \leq m$ where G_1 (resp. G_2) is the matrix representation group of the action of \widetilde{H} on the left-hand side $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a_i}) \oplus F^{\oplus b_1}$ (resp. the right-hand side $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a'_i}) \oplus F'^{\oplus b'_1}$) of the isomorphism (8) and $m = \text{rank}_{\mathbb{Z}} \text{Hom}(G_1, G_2)$.

However, in general, we have that $\det P_i \neq \pm 1$ for any $1 \leq i \leq m$. In the general case, we should seek a matrix P with $\det P = \pm 1$ which is given as a linear combination $P = \sum_{i=1}^m c_i P_i$. This task is important for us and not easy in general even if we use a computer.

We made the following GAP algorithms which may find a matrix $P = \sum_{i=1}^m c_i P_i$ with $G_1 P = P G_2$ and $\det P = \pm 1$.

We will explain the algorithms below when the input \mathcal{P} is obtained by StablyEquivalentFCheckPSubdirectProduct(\widetilde{H} , l_1, l_2) although it works in more general situations.

SearchPRowBlocks

▸ SearchPRowBlocks(P)

returns the records bpBlocks and rowBlocks where bpBlocks (resp. rowBlocks) is the decomposition of the list $l = [1, \dots, m]$ (resp. $l = [1, \dots, n]$) with $m = \text{rank}_{\mathbb{Z}} \text{Hom}(G_1, G_2)$ (resp. $n = \text{size } G_1$) according to the direct sum decomposition of M_{G_1} for a basis $\mathcal{P} = \{P_1, \dots, P_m\}$ of the solution space of $G_1 P = P G_2$ where G_1 (resp. G_2) is the matrix representation group of the action of \widetilde{H} on the left-hand side $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a_i}) \oplus F^{\oplus b_1}$ (resp. the right-

hand side $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a'_i}) \oplus F' \oplus b'_1)$ of the isomorphism (8).

We write $B[t] = \text{SearchPRowBlocks}(\mathcal{P}).\text{bpBlocks}[t]$ and $R[t] = \text{SearchPRowBlocks}(\mathcal{P}).\text{rowBlocks}[t]$.

SearchPFilterRowBlocks

```
▸ SearchPFilterRowBlocks( $P, B[t], R[t], j$ )
```

returns the lists $\{M_s\}$ where M_s is the $n_t \times n$ matrix with all invariant factors 1 which is of the form $M_s = \sum_{i \in B[t]} c_i P'_i$ ($c_i \in \{0, 1\}$) at most j non-zero c_i 's and P'_i is the submatrix of P_i consists of $R[t]$ rows with $n_t = \text{length}(R[t])$ for a basis $\mathcal{P} = \{P_1, \dots, P_m\}$ of the solution space of $G_1 P = P G_2$ where G_1 (resp. G_2) is the matrix representation group of the action of \widetilde{H} on the left-hand side $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a_i}) \oplus F^{\oplus b_1}$ (resp. the right-hand side $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a'_i}) \oplus F' \oplus b'_1)$ of the isomorphism (8), $B[t] = \text{SearchPRowBlocks}(\mathcal{P}).\text{bpBlocks}[t]$, $R[t] = \text{SearchPRowBlocks}(\mathcal{P}).\text{rowBlocks}[t]$ and $j \geq 1$.

```
▸ SearchPFilterRowBlocks( $P, B[t], R[t], j, C$ )
```

returns the same as $\text{SearchPFilterRowBlocks}(P, B[t], R[t], j)$ but with respect to $c_i \in C$ instead of $c_i \in \{0, 1\}$ for the list C of integers.

SearchPFilterRowBlocksRandomMT

```
▸ SearchPFilterRowBlocksRandomMT( $P, B[t], R[t], u$ )
```

returns the same as $\text{SearchPFilterRowBlocks}(P, B[t], R[t], j)$ but with respect to random u c_i 's via Mersenne Twister instead of at most j non-zero c_i 's for integer $u \geq 1$.

```
▸ SearchPFilterRowBlocksRandomMT( $P, B[t], R[t], u, C$ )
```

returns the same as $\text{SearchPFilterRowBlocksRandomMT}(P, B[t], R[t], u)$ but with respect to $c_i \in C$ instead of $c_i \in \{0, 1\}$ for the list C of integers.

SearchPMergeRowBlock

```
▸ SearchPMergeRowBlock( $m1, m2$ )
```

returns all concatenations of the matrices M_s and M_t vertically with all invariant factors 1 (resp. a concatenation of the matrices M_s and M_t vertically with determinant ± 1) for $m_1 = \{M_s\}$ and $m_2 = \{M_t\}$ where M_s are $n_1 \times n$ matrices and M_t are $n_2 \times n$ matrices with $n_1 + n_2 < n$ (resp. $n_1 + n_2 = n$).

When there exists $t \in \mathbb{Z}$ such that $R[t] = \{j\}$, we can use:

SearchPLinear

▸ SearchPLinear($M, P1$)

returns the list $\{\det(M + P_i)\}_{i \in B[t]}$ of integers for an $n \times n$ matrix M which is obtained by inserting the zero row into the j -th row of $(n - 1) \times n$ matrix $M_s = \sum_{i \notin B[t]} c_i P'_i$ with all invariant factors 1 and $\mathcal{P}_1 = \{P_i\}_{i \in B[t]}$ where $B[t] = \text{SearchPRowBlocks}(\mathcal{P}).\text{bpBlocks}[t]$, P'_i is the submatrix of P_i deleting the j -th row, and $\mathcal{P} = \{P_1, \dots, P_m\}$ is obtained by $\text{StablyEquivalentFCheckPSubdirectProduct}(\widetilde{H}, l_1, l_2)$ under the assumption that there exists $t \in \mathbb{Z}$ such that $R[t] = \{j\}$.

When there exist $t_1, t_2 \in \mathbb{Z}$ such that $R[t_1] = \{j_1\}$, $R[t_2] = \{j_2\}$, we can use:

SearchPBilinear

▸ SearchPBilinear($M, P1, P2$)

returns the matrix $[\det(M + P_{i_1} + P_{i_2})]_{i_1 \in B[t_1], i_2 \in B[t_2]}$ for an $n \times n$ matrix M which is obtained by inserting the two zero rows into the j_1 -th row and the j_2 -th row of $(n - 2) \times n$ matrix $M_s = \sum_{i \notin B[t_1] \cup B[t_2]} c_i P'_i$ with all invariant factors 1 and $\mathcal{P}_1 = \{P_{i_1}\}_{i_1 \in B[t_1]}$, $\mathcal{P}_2 = \{P_{i_2}\}_{i_2 \in B[t_2]}$, where $B[t_1] = \text{SearchPRowBlocks}(\mathcal{P}).\text{bpBlocks}[t_1]$, $B[t_2] = \text{SearchPRowBlocks}(\mathcal{P}).\text{bpBlocks}[t_2]$, P'_i is the submatrix of P_i deleting the j_1 -th and the j_2 -th rows, and $\mathcal{P} = \{P_1, \dots, P_m\}$ is obtained by $\text{StablyEquivalentFCheckPSubdirectProduct}(\widetilde{H}, l_1, l_2)$ under the assumption that there exist $t_1, t_2 \in \mathbb{Z}$ such that $R[t_1] = \{j_1\}$ and $R[t_2] = \{j_2\}$.

When there exists $t \in \mathbb{Z}$ such that $R[t] = \{j_1, j_2\}$, we can use:

SearchPQuadratic

▸ SearchPQuadratic($M, P1$)

returns the matrix $[\frac{1}{2}(\det(M + P_{i_1} + P_{i_2}) - \det(M + P_{i_1}) - \det(M + P_{i_2}))]_{i_1, i_2 \in B[t]}$ for an $n \times n$ matrix M which is obtained by inserting the two zero rows into the j_1 -th row and the j_2 -th row of $(n - 2) \times n$ matrix $M_s = \sum_{i \notin B[t]} c_i P'_i$ with all invariant factors 1 and $\mathcal{P}_1 = \{P_i\}_{i \in B[t]}$, where $B[t] = \text{SearchPRowBlocks}(\mathcal{P}).\text{bpBlocks}[t]$, P'_i is the submatrix of P_i deleting the j_1 -th and j_2 -th rows and $\mathcal{P} = \{P_1, \dots, P_m\}$ is obtained by $\text{StablyEquivalentFCheckPSubdirectProduct}(\widetilde{H}, l_1, l_2)$ under the assumption that there exists $t \in \mathbb{Z}$ such that $R[t] = \{j_1, j_2\}$.

When $R[1] = \{1, \dots, m\}$, we can use:

SearchP1

▸ SearchP1(P)

returns a matrix $P = \sum_{i=1}^m c_i P_i$ with $c_i \in \{0, 1\}$, $G_1 P = P G_2$ and $\det P = \pm 1$ where G_1 (resp. G_2) is the matrix representation group of the action of \widetilde{H} on the left-hand side $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a_i}) \oplus F^{\oplus b_1}$ (resp. the right-hand side $(\oplus_{i=1}^r \mathbb{Z}[\widetilde{H}/H_i]^{\oplus a'_i}) \oplus F'^{\oplus b'_1}$) of the isomorphism (8) for $\mathcal{P} = \{P_1, \dots, P_m\}$ which is obtained by

StablyEquivalentFCheckPSubdirectProduct(\widetilde{H}, l_1, l_2) under the assumption that $R[1] = \{1, \dots, m\}$.

▸ SearchP1(P, C)

returns the same as SearchP1(P) but with respect to $c_i \in C$ instead of $c_i \in \{0, 1\}$ for the list C of integers.

Endomorphismring

▸ Endomorphismring(G)

returns a \mathbb{Z} -basis of $\text{End}_{\mathbb{Z}[G]}(M_G)$ for a finite subgroup G of $\text{GL}(n, \mathbb{Z})$.

IsCodimJacobsonEnd1

▸ IsCodimJacobsonEnd1(G, p)

returns true (resp. false) if $\dim_{\mathbb{Z}/p\mathbb{Z}}(E/pE)/J(E/pE) = 1$ (resp. $\neq 1$) where $E = \text{End}_{\mathbb{Z}[G]}(M_G)$ for a finite subgroup G of $\text{GL}(n, \mathbb{Z})$ and prime number p . If this returns true, then $M_G \otimes_{\mathbb{Z}} \mathbb{Z}_p$ is an indecomposable $\mathbb{Z}_p[G]$ -lattice. In particular, M_G is an indecomposable G -lattice (see [HY, Lemma 6.11]).

IdempotentsModp

▸ IdempotentsModp(B, p)

returns all idempotents of R/pR for a \mathbb{Z} -basis B of a subring R of $n \times n$ matrices $M(n, \mathbb{Z})$ over \mathbb{Z} and prime number p . If this returns only the zero and the identity matrices when $R = \text{End}_{\mathbb{Z}[G]}(M_G)$, then $M_G \otimes_{\mathbb{Z}} \mathbb{Z}_p$ is an indecomposable $\mathbb{Z}_p[G]$ -lattice. In particular, M_G is an indecomposable G -lattice (see [HY, Lemma 6.10]).

ConjugacyClassesSubgroups2WSEC

▸ `ConjugacyClassesSubgroups2WSEC(G)`

returns the records `ConjugacyClassesSubgroups2` and `WSEC` where `ConjugacyClassesSubgroups2` is the list $[g_1, \dots, g_m]$ of conjugacy classes of subgroups of $G \leq \text{GL}(n, \mathbb{Z})$ ($n = 3, 4$) with the fixed ordering via the function `ConjugacyClassesSubgroups2(G)` (see [HY17, Section 4.1]) and `WSEC` is the list $[w_1, \dots, w_m]$ where g_i is in the w_i -th weak stably k -equivalent class WSEC_{w_i} in dimension n .

MaximalInvariantNormalSubgroup

▸ `MaximalInvariantNormalSubgroup(G , ConjugacyClassesSubgroups2WSEC(G))`

returns a maximal normal subgroup N of G which satisfies that $\pi(H_1) = \pi(H_2)$ implies $\psi(H_1) = \psi(H_2)$ for any $H_1, H_2 \leq G$ where $\pi : G \rightarrow G/N$ is the natural homomorphism, $\psi : H_i \mapsto w_i$, and H_i is in the w_i -th weak stably k -equivalent class WSEC_{w_i} in dimension n .

PossibilityOfStablyEquivalentSubdirectProducts with "WSEC" option

▸ `PossibilityOfStablyEquivalentSubdirectProducts(G, G' , ConjugacyClassesSubgroups2WSEC(G), ConjugacyClassesSubgroups2WSEC(G'), ["WSEC"])`

returns the list l of the subdirect products $\widetilde{H} \leq G \times G'$ of G and G' up to $(\text{GL}(n_1, \mathbb{Z}) \times \text{GL}(n_2, \mathbb{Z}))$ -conjugacy which satisfy $w_1 = w_2$ for any $H \leq \widetilde{H}$ where $\varphi_i(H)$ is in the w_i -th weak stably k -equivalent class WSEC_{w_i} in dimension n ($n = 3, 4$) and $\widetilde{H} \leq G \times G'$ is a subdirect product of G and G' which acts on M_G and $M_{G'}$ through the surjections $\varphi_1 : \widetilde{H} \rightarrow G$ and $\varphi_2 : \widetilde{H} \rightarrow G'$ respectively (indeed, this function computes it for H up to conjugacy for the sake of saving time).

IsomorphismFromSubdirectProduct

▸ `IsomorphismFromSubdirectProduct($H \sim$)`

returns the isomorphism $\sigma : G/N \rightarrow G'/N'$ which satisfies $\sigma(\varphi_1(h)N) = \varphi_2(h)N'$ for any $h \in \widetilde{H}$ where $N = \varphi_1(\text{Ker}(\varphi_2))$ and $N' = \varphi_2(\text{Ker}(\varphi_1))$ for a subdirect product $\widetilde{H} \leq G \times G'$ of G and G' with surjections $\varphi_1 : \widetilde{H} \rightarrow G$ and $\varphi_2 : \widetilde{H} \rightarrow G'$.

AutGSubdirectProductsWSECInvariant

▸ `AutGSubdirectProductsWSECInvariant(G)`

returns subdirect products $\widetilde{H}_m = \{(g, g^{\sigma_m}) \mid g \in G, g^{\sigma_m} \in G^{\sigma_m}\}$

$(1 \leq m \leq s)$ of G and G^{σ_m} where $\{\sigma_1, \dots, \sigma_s\}$ is a complete set of representatives of the double coset $X \backslash Z / X$,

$$\text{Inn}(G) \leq X \leq Y \leq Z \leq \text{Aut}(G),$$

$$X = \text{Aut}_{\text{GL}(n, \mathbb{Z})}(G) = \{\sigma \in \text{Aut}(G) \mid \text{there exists } u \in \text{GL}(n, \mathbb{Z}) \text{ such that } u^{-1} \sigma u \simeq N_{\text{GL}(n, \mathbb{Z})}(G) / Z_{\text{GL}(n, \mathbb{Z})}(G),$$

$$Y = \{\sigma \in \text{Aut}(G) \mid [M_G]^{fl} = [M_{G^\sigma}]^{fl} \text{ as } \widetilde{H}\text{-lattices where } \widetilde{H} = \{(g, g^\sigma) \mid g \in G\},$$

$$Z = \{\sigma \in \text{Aut}(G) \mid [M_H]^{fl} \sim [M_{H^\sigma}]^{fl} \text{ for any } H \leq G\},$$

$\text{Inn}(G)$ is the group of inner automorphisms on G , $\text{Aut}(G)$ is the group of automorphisms on G , $N_{\text{GL}(n, \mathbb{Z})}(G)$ is the normalizer of G in $\text{GL}(n, \mathbb{Z})$ and $Z_{\text{GL}(n, \mathbb{Z})}(G)$ is the centralizer of G in $\text{GL}(n, \mathbb{Z})$.

AutGSubdirectProductsWSECIInvariantGen

▸ `AutGSubdirectProductsWSECIInvariantGen(G)`

returns the same as `AutGSubdirectProductsWSECIInvariant(G)` but with respect to $\{\sigma_1, \dots, \sigma_t\}$ where $\sigma_1, \dots, \sigma_t \in Z$ are some minimal number of generators of the double cosets of $X \backslash Z / X$, i.e. minimal number of elements $\sigma_1, \dots, \sigma_t \in Z$ which satisfy $\langle \sigma_1, \dots, \sigma_t, x \mid x \in X \rangle = Z$, instead of a complete set of representatives of the double coset $X \backslash Z / X$. If this returns `[]`, then we get $X = Y = Z$.

AutGLnZ

▸ `AutGLnZ(G)`

returns

$$X = \text{Aut}_{\text{GL}(n, \mathbb{Z})}(G) = \{\sigma \in \text{Aut}(G) \mid \text{there exists } u \in \text{GL}(n, \mathbb{Z}) \text{ such that } u^{-1} \sigma u \simeq N_{\text{GL}(n, \mathbb{Z})}(G) / Z_{\text{GL}(n, \mathbb{Z})}(G).$$

N3WSECMembersTable

▸ `N3WSECMembersTable[r][i]`

returns an integer j which satisfies that $N_{3,j}$ is the i -th group in the weak stably k -equivalent class WSEC_r .

N4WSECMembersTable

▸ `N4WSECMembersTable[r][i]`

is the same as `N3WSECMembersTable[r][i]` but using $N_{4,j}$ instead of $N_{3,j}$.

I4WSECMembersTable

▸ I4WSECMembersTable[*r*][*i*]

is the same as N3WSECMembersTable[*r*][*l*] but using $I_{4,j}$ instead of $N_{3,j}$.

AutGWSECIInvariantSmallDegreeTest

▸ AutGWSECIInvariantSmallDegreeTest(*G*)

returns the list $l = [l_1, \dots, l_s]$ ($l_1 \leq \dots \leq l_s$) of integers with the minimal l_s, \dots, l_1 which satisfies $Z = Z'$ where

$$Z = \{\sigma \in \text{Aut}(G) \mid [M_H]^{fl} \sim [M_{H^\sigma}]^{fl} \text{ for any } H \leq G\},$$

$$Z' = \{\sigma \in \text{Aut}(G) \mid [M_H]^{fl} \sim [M_{H^\sigma}]^{fl} \text{ for any } H \leq G \text{ with } [G : H] \in l\}$$

for $G \leq \text{GL}(n, \mathbb{Z})$ ($n = 3, 4$).

MaximalInvariantNormalSubgroups

▸ MaximalInvariantNormalSubgroups(*G*, ConjugacyClassesSubgroups2WSEC(*G*))

returns maximal normal subgroups N of G which satisfy that $\pi(H_1) = \pi(H_2)$ implies $\psi(H_1) = \psi(H_2)$ for any $H_1, H_2 \leq G$ where $\pi : G \rightarrow G/N$ is the natural homomorphism, $\psi : H_i \mapsto w_i$, and H_i is in the w_i -th weak stably k -equivalent class WSEC_{w_i} in dimension n . If this returns [Group([])], then $N = 1$ where $\widetilde{H} \leq G \times G$ is a subdirect product of G and G with surjections $\varphi_1 : \widetilde{H} \rightarrow G$, $\varphi_2 : \widetilde{H} \rightarrow G$ and $N = \varphi_1(\text{Ker}(\varphi_2)) \triangleleft G$.

AutWSEC

▸ AutWSEC(*G*)

returns

$$Z = \{\sigma \in \text{Aut}(G) \mid [M_H]^{fl} \sim [M_{H^\sigma}]^{fl} \text{ for any } H \leq G\}.$$

IdCoset

▸ IdCoset(*G*, *H*)

returns $[d, m]$ when the action of G on G/H may be regarded as $G \simeq dTm \leq S_d$ with $d = [G : H] > 1$. If $d = 1$, this returns [1].

References

[HHY20] Sumito Hasegawa, Akinari Hoshi and Aiichi Yamasaki, Rationality

problem for norm one tori in small dimensions, Math. Comp. **89** (2020) 923-940. [AMS](#) Extended version: [arXiv:1811.02145](#).
[HY17] Akinari Hoshi and Aiichi Yamasaki, Rationality problem for algebraic tori, Mem. Amer. Math. Soc. **248** (2017) no. 1176, v+215 pp. [AMS](#) Preprint version: [arXiv:1210.4525](#).
[HY] Akinari Hoshi and Aiichi Yamasaki, Birational classification for algebraic tori, [arXiv:2112.02280](#).

Copyright © 2021 Akinari Hoshi, Aiichi Yamasaki